

Refine Search

Search Results -

Term	Documents
WESLOARC	1
WESLOARCS	0
(WESLOARC AND 1).PGPB,USPT.	1
(L1 AND WESLOARC).PGPB,USPT.	1

Database:

US Pre-Grant Publication Full-Text Database
 US Patents Full-Text Database
 US OCR Full-Text Database
 EPO Abstracts Database
 JPO Abstracts Database
 Derwent World Patents Index
 IBM Technical Disclosure Bulletins

Search:

L2

Refine Search

Recall Text

Clear

Interrupt

Search History

 DATE: Monday, December 13, 2004 [Printable Copy](#) [Create Case](#)

<u>Set</u> <u>Name</u> side by side	<u>Query</u>	<u>Hit</u> <u>Count</u>	<u>Set</u> <u>Name</u> result set
	<i>DB=PGPB,USPT; THES=ASSIGNEE; PLUR=YES; OP=ADJ</i>		
<u>L2</u>	L1 and wesloarc	1	<u>L2</u>
<u>L1</u>	(lobelia or lobelias or lobelia adj X adj hybrida or lobelia adj X adj hybrids or lobelia adj X adj hybrid or lobelia adj erinus)	150	<u>L1</u>

END OF SEARCH HISTORY

Hit List

Clear	Generate Collection	Print	Fwd Refs	Bkwd Refs
Generate OACS				

Search Results - Record(s) 1 through 1 of 1 returned.

☐ 1. Document ID: US 20040199971 P1

Using default format because multiple data bases are involved.

L2: Entry 1 of 1

File: PGPB

Oct 7, 2004

PGPUB-DOCUMENT-NUMBER: 20040199971

PGPUB-FILING-TYPE: new

DOCUMENT-IDENTIFIER: US 20040199971 P1

TITLE: Lobelia plant named 'Weslospot'

PUBLICATION-DATE: October 7, 2004

INVENTOR-INFORMATION:

NAME	CITY	STATE	COUNTRY	RULE-47
Westhoff, Heinrich	Sudlohn		DE	

US-CL-CURRENT: PLT/268

FULL	Title	Citation	Front	Review	Classification	Date	Reference	Sequences	Attachments	Claims	KUUC	Drawn De
------	-------	----------	-------	--------	----------------	------	-----------	-----------	-------------	--------	------	----------

Clear	Generate Collection	Print	Fwd Refs	Bkwd Refs	Generate OACS
-------	---------------------	-------	----------	-----------	---------------

Term	Documents
WESLOARC	1
WESLOARCS	0
(WESLOARC AND 1).PGPB,USPT.	1
(L1 AND WESLOARC).PGPB,USPT.	1

Display Format:

[Previous Page](#)

[Next Page](#)

[Go to Doc#](#)

Set Items Description

diology 12/13/04

?s lobelia or lobelias or lobelia X hybrids or lobelia X hybrida or lobelia X hybrid or
lobelia hybrid or lobelia hybrids or lobelia hybrida

2350 LOBELIA

98 LOBELIAS

0 LOBELIA X HYBRIDS

0 LOBELIA X HYBRIDA

0 LOBELIA X HYBRID

0 LOBELIA HYBRID

0 LOBELIA HYBRIDS

1 LOBELIA HYBRIDA

S1 2388 LOBELIA OR LOBELIAS OR LOBELIA X HYBRIDS OR LOBELIA X
HYBRIDA OR LOBELIA X HYBRID OR LOBELIA HYBRID OR LOBELIA
HYBRIDS OR LOBELIA HYBRIDA

?s s1 and wesloarc

2388 S1

1 WESLOARC

S2 1 S1 AND WESLOARC

?

2/9,K/1 (Item 1 from file: 654)
DIALOG(R) File 654:US Pat.Full.
(c) Format only 2004 The Dialog Corp. All rts. reserv.

0005827559

Lobelia plant named 'Weslospot'
Inventor: Westhoff, Heinrich, INV
Assignee: Josef und Heinrich Westhoff Gartenbau- Spezialkulturen (03),
Sudlohn, DE
Correspondence Address: Catherine Anne Whealy Proprietary Rights
International, P.O. Box 220, Tolar, TX, 76476, US

	Publication Number	Kind	Date	Application Number	Filing Date
Main Patent	US 20040199971	P1	20041007	US 2004805581	20040320
Priority				EP 2003198	20030203

US Classification on document (Main): PLT268000
International Classification (Edition 07): A01H-005/00

Fulltext Word Count: 1993
Number of Claims: 1
Exemplary or Independent Claim Number(s): 1
Number of Drawing Sheets: 1
Number of Figures: 2

Abstract:

A new and distinct cultivar of **Lobelia** plant named 'Weslospot', characterized by its cascading plant habit; freely branching habit with short internodes; continuously and freely flowering habit; densely pubescent leaves; light blue and white bi-colored flowers; and relatively tolerant to high temperatures

Summary of the Invention:

BOTANICAL CLASSIFICATION/CULTIVAR DESIGNATION

[0001] **Lobelia**Xhybrida cultivar WESLOSPOT

BACKGROUND OF THE INVENTION

[0002] The present Invention relates to a new and distinct cultivar of **Lobelia** plant, botanically known as **Lobelia** erinus, and hereinafter referred to by the name 'Weslospot'.

[0003] The new **Lobelia** is a product of a planned breeding program conducted by the Inventor in S[u, dieresis or umlaut mark]dlohn, Germany. The objective of the breeding program was to develop new **Lobelia** cultivars with continuous flowering, interesting flower colors and high temperature tolerance.

[0004] The new **Lobelia** originated from a cross-pollination made by the Inventor in 2002 of a proprietary selection of **Lobelia** erinus identified as code number 01P050, not patented, as the female, or seed, parent with a proprietary selection of **Lobelia** erinus identified as code number 01P508, not patented, as the male, or pollen, parent. The new **Lobelia** was discovered and selected by the Inventor from within the resultant progeny from the above-mentioned cross-pollination in a controlled environment in S[u, dieresis or umlaut mark]dlohn, Germany in 2002.

[0005] Asexual reproduction since 2002 of the new cultivar by terminal cuttings in a controlled environment in S[u, dieresis or umlaut mark]dlohn, Germany, has shown that the unique features of this new **Lobelia** are stable and reproduced true to type in successive generations

SUMMARY OF THE INVENTION

[0006] The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Weslospot'. These characteristics in combination distinguish 'Weslospot' as a new and distinct cultivar:

- [0007] 1. Cascading plant habit.
- [0008] 2. Freely branching habit with short internodes.
- [0009] 3. Continuously and freely flowering habit.
- [0010] 4. Densely pubescent leaves.
- [0011] 5. Light blue and white bi-colored flowers.
- [0012] 6. Relatively tolerant to high temperatures.

[0013] Plants of the new **Lobelia** have larger flowers than plants of the female parent, a proprietary selection identified as code number 01P050. In addition, plants of the new **Lobelia** and the female parent selection differ in flower color. Plants of the new **Lobelia** have foliage with dense pubescence compared to the smooth foliage of plants of the male parent, a proprietary selection identified as code number 01P508. In addition, plants of the new **Lobelia** have plants with thicker stems and lighter colored flowers than plants of the male parent selection.

[0014] Plants of the new **Lobelia** differ primarily from plants of the cultivar **Wesloarc**, disclosed in a U.S. Plant patent application filed concurrently, primarily in flower color.

[0015] Plants of the cultivar Weslospot can be compared to the cultivar Weslowei, disclosed in U.S. Plant Pat. No. 12,708. However, in side-by-side comparisons conducted in S[u, dieresis or umlaut mark]dlohn, Germany, plants of the cultivar Weslospot and the cultivar Weslowei differed in the following characteristics:

- [0016] 1. Leaves and stems of plants of the new **Lobelia** were more pubescent than leaves and stems of plants of the cultivar Weslowei.
- [0017] 2. Plants of the new **Lobelia** flowered more continuously than plants of the cultivar Weslowei.
- [0018] 3. Plants of the new **Lobelia** and the cultivar Weslowei differed in flower color.
- [0019] 4. Sepals of plants of the new **Lobelia** were smaller than sepals of plants of the cultivar Weslowei.
- [0020] 5. Plants of the new **Lobelia** were more tolerant to high temperatures than plants of the cultivar Weslowei.

[0021] Plants of the new **Lobelia** can also be compared to plants of the cultivar Weslobigblue, disclosed in U.S. Plant Pat. No. 12,634. However, in side-by-side comparisons conducted in S[u, dieresis or umlaut mark]dlohn, Germany, plants of the new **Lobelia** differed from plants of the cultivar Weslobigblue in the following characteristics:

- [0022] 1. Leaves and stems of plants of the new **Lobelia** were more pubescent and differed in color compared to leaves and stems of plants of the cultivar Weslobigblue.
- [0023] 2. Stems of plants of the new **Lobelia** were shorter and narrower than stems of plants of the cultivar Weslobigblue.
- [0024] 3. Plants of the new **Lobelia** had smaller flowers than plants of the cultivar Weslobigblue.
- [0025] 4. Flowers of plants of the new **Lobelia** were lighter blue in color than flowers of plants of the cultivar Weslobigblue.

Description of the Drawings:

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

[0026] The accompanying colored photographs illustrate the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description, which accurately describe the actual colors of the new **Lobelia**. The photograph at the top of the sheet comprises a side view of a typical plant of 'Weslospot' grown in a window box container. The photograph at the bottom of the sheet comprises a close-up view of a typical individual flower of 'Weslospot'.

Description of the Invention:

DETAILED BOTANICAL DESCRIPTION

[0027] Plants of the cultivar Weslospot have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity without, however, any variance in genotype.

[0028] In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2001 Edition, except where general terms of ordinary dictionary significance are used. Plants used for the description were grown in a glass-covered greenhouse and conditions that closely approximate commercial production conditions during the spring and summer in S[u, dieresis or umlaut mark]dlohn, Germany. Plants used for the above-mentioned photographs and following description were grown as one plant per 12-cm container or three plants per 25-cm hanging basket container. During the production of the plants, day temperatures ranged from 20 to 25 degree C. and night temperatures ranged from 1 to 18 degree C. Plants were pinched once during the production period by removing about 1 to 2 cm of the uppermost apical growing tip. Plants were about 20 weeks from planting when the photographs and description were taken.

BOTANICAL CLASSIFICATION

[0029] **Lobelia** erinus cultivar Weslospot.

PARENTAGE

[0030] Female parent: Proprietary selection of **Lobelia** erinus identified as code number 01P050, not patented [0031] Male parent: Proprietary selection of **Lobelia** erinus identified as code number 01P508, not patented.

PROPAGATION

[0032] Type cutting: Terminal vegetative cuttings.

[0033] Time to initiate roots: About 18 to 21 days at 20 degree C.

[0034] Time to develop roots: About 20 to 28 days at 20 degree C [0035] Root description: Fine, fibrous and well-branched.

PLANT DESCRIPTION

[0036] Plant form/habit: Cascading flowering plants with light blue and white-colored flowers. Lateral shoots outwardly spreading; plants uniform with dense foliage. Foliage and stems are very pubescent. Freely branching with lateral branches forming at every node; dense and bushy plant habit. Pinching plants enhances branching. Moderately vigorous growth habit.

[0037] Usage: Appropriate for hanging baskets, window boxes and patio containers.

[0038] Plant height (soil level to top of plant plane): About 16 cm.

[0039] Plant length (soil level to lateral branches apices): About 68 cm.

[0040] Plant diameter: About 50 to 60 cm.

[0041] Lateral branch description:

[0042]Length.- About 17 cm.

[0043]Diameter.- About 1.6 mm.

[0044]Internode length.- About 1.7 cm.

[0045]Texture.- Densely pubescent.

[0046]Color.- 138A.

[0047] Foliage description:

[0048]Arrangement.- Alternate; simple.

[0049]Basal leaves.- Length: About 4.8 cm. Width: About 2.9 cm. Shape: Nearly round. Apex: Retuse. Base: Attenuate. Margin: Slightly crenate. Petiole length: About 1.5 mm.

[0050]Mid -plant leaves.-Length: About 3.5 cm. Width: About 2.1 cm. Shape: Ovate. Apex: Rounded. Base: Attenuate. Margin: Crenate. Petiole length: About 1 cm.

[0051]Apical leaves.- Length: About 2.7 cm. Width: About 7.7 mm. Shape: Oblanceolate. Apex: Slightly acute. Base: Attenuate. Margin: Mostly entire. Petiole length: Petioles not observed.

[0052]Texture, all leaves, upper and lower surfaces.- Densely pubescent.

[0053]Color, all leaves.- Developing foliage, upper surface: 146A. Developing foliage, lower surface: 146B. Fully developed foliage, upper surface: 147A; venation, 147B. Fully developed foliage, lower surface: 147B; venation, 147B.

FLOWER DESCRIPTION

[0054] Flower type and habit: Flowers arranged singly at lateral apices. Flowers held mostly outwardly. Flowers persistent. Older flowers are overgrown by new flowers and foliage. Freely and continuously flowering. Flowers not fragrant.

[0055] Flower shape: Tubular with three larger lower petals and two upright petals.

[0056] Natural flowering season: Spring until frost in the autumn.

[0057] Flower longevity on the plant: Longevity of individual flowers is highly dependent on weather conditions; typically three to ten days.

[0058] Flower size:

[0059]Diameter.- About 1.9 cm.

[0060]Depth (height).-About 1.8 cm.

[0061]Tube length.- About 1 cm.

[0062]Throat diameter, distal end.- About 4.1 mm.

[0063]Tube diameter, proximal end.- About 2.7 mm.

[0064] Flower buds:

[0065]Length.- About 1.1 cm.

[0066]Diameter.- About 2.4 mm.

[0067]Shape.- Oblong.

[0068]Color.- Towards the base, 145D; towards the apex, close to 145C; with development, color becoming closer to 122B to 122C.

[0069] Petals:

[0070]Arrangement.- Single whorl of five petals, fused; three larger lower petals and two smaller upper petals.

[0071]Three lower petals.- Shape: Obovate. Length, above throat: About 1.2 cm. Width: About 6.4 mm.

[0072]Two upper petals.- Shape: Spatulate. Length, above throat: About 6.7 mm. Width: About 3.1 mm.

[0073]Upper and lower petals.- Apex: Cuspidate. Margin: Entire. Texture, upper and lower surfaces: Smooth, satiny. Color: When opening, upper surface: 96D, towards the center, 155C. When opening, lower surface: 97B, towards the center, 155C. Fully opened, upper surface: 97A to 97B, towards the center, 155C. Petal, fully opened, lower surface: 97B, towards the center, 155C. Throat: 155C; spots, 93A to 93C. Tube: 97C to 97D; spots, 97B.

[0074] Sepals:

[0075]Arrangement.- Single whorl of five sepals, star-shaped calyx.

[0076]Length.- About 7.7 mm.

[0077]Width.- About 1.1 mm.

[0078]Shape.- Elliptic.

[0079]Apex.- Acute.

[0080]Margin.- Entire.

[0081]Texture, upper and lower surfaces.- Smooth.

[0082]Color, upper and lower surfaces.- 137A.

[0083] Peduncles:

[0084]Appearance.- Wiry.

[0085]Length.- About 2.3 cm.

[0086]Diameter.- About 9 mm.

[0087]Texture.- Densely pubescent.

[0088]Color.- 137A.

[0089] Reproductive organs:

[0090]Stamens.- Quantity per flower: About five, fused. Anther length: About 2.5 mm. Anther diameter: About 1.8 mm. Anther color: 186A. Pollen amount: Moderate. Pollen color: 8A.

[0091] Pistils.- Quantity per flower: One. Pistil length: About 6.2 mm. Stigma shape: Two - parted, ovate. Stigma texture: Pubescent. Stigma color, immature: 186A. Stigma color, fully mature: 81A to 81C. Style length: About 4 mm. Style color: 145A. Ovary color: 145A.

[0092] Seed/fruit.- Seed and fruit production have not been observed.

DISEASE/PEST RESISTANCE

[0093] Plants of the new **Lobelia** have not been noted to be resistant to pathogens and pests common to **Lobelia**.

TEMPERATURE TOLERANCE

[0094] Plants of the new **Lobelia** have been observed to tolerate temperatures ranging from 2 to 30 degree C.

It is claimed:

Exemplary or Independent Claim(s):

1. A new and distinct cultivar of **Lobelia** plant named 'Weslospot', as illustrated and described.

Lobelia plant named 'Weslospot'

Abstract:

A new and distinct cultivar of **Lobelia** plant named 'Weslospot', characterized by its cascading plant habit; freely branching habit with short internodes...

Summary of the Invention:

...0002] The present Invention relates to a new and distinct cultivar of **Lobelia** plant, botanically known as **Lobelia** erinus, and hereinafter referred to by the name 'Weslospot'...

...0003] The new **Lobelia** is a product of a planned breeding ...or umlaut mark]dlohn, Germany. The objective of the breeding program was to develop new **Lobelia** cultivars with continuous flowering, interesting flower colors and high temperature tolerance...

...0004] The new **Lobelia** originated from a cross-pollination made by the Inventor in 2002 of a proprietary selection of **Lobelia** erinus identified as code number 01P050, not patented, as the female, or seed, parent with a proprietary selection of **Lobelia** erinus identified as code number 01P508, not patented, as the male, or pollen, parent. The new **Lobelia** was discovered and selected by the Inventor from within the resultant progeny from the above...

...dieresis or umlaut mark]dlohn, Germany, has shown that the unique features of this new **Lobelia** are stable and reproduced true to type in successive generations...0013] Plants of the new **Lobelia** have larger flowers than plants of the female parent, a proprietary selection identified as code number 01P050. In addition, plants of the new **Lobelia** and the female parent selection differ in flower color. Plants of the new **Lobelia** have foliage with dense pubescence compared to the smooth foliage of plants of the male parent, a proprietary selection identified as code number 01P508. In addition, plants of the new **Lobelia** have plants with thicker stems and lighter colored flowers than plants of the male parent...0014] Plants of the new **Lobelia** differ primarily from plants of the cultivar **Wesloarc**, disclosed in a U.S. Plant patent application filed concurrently, primarily in flower color...

...0016] 1. Leaves and stems of plants of the new **Lobelia** were more pubescent than leaves and stems of plants of the cultivar **Weslowei**...

...0017] 2. Plants of the new **Lobelia** flowered ...0018] 3. Plants of the new **Lobelia** and the cultivar **Weslowei** differed in flower color...

...0019] 4. Sepals of plants of the new **Lobelia** were smaller than sepals of plants of the cultivar Weslowei...

...0020] 5. Plants of the new **Lobelia** were more tolerant to high temperatures than plants of the cultivar Weslowei...

...0021] Plants of the new **Lobelia** can also be compared to plants of the cultivar Weslobigblue, disclosed in U.S. Plant dlohn, Germany, plants of the new **Lobelia** differed from plants of the cultivar Weslobigblue in the following characteristics...

...0022] 1. Leaves and stems of plants of the new **Lobelia** were more pubescent and differed in color compared to leaves and stems of plants of ...

...0023] 2. Stems of plants of the new **Lobelia** were shorter and narrower than stems of plants of the cultivar Weslobigblue...

...0024] 3. Plants of the new **Lobelia** had smaller flowers than plants of the cultivar Weslobigblue...

...0025] 4. Flowers of plants of the new **Lobelia** were lighter blue in color than flowers of

Description of the Drawings:

...cited in the detailed botanical description, which accurately describe the actual colors of the new **Lobelia**. The photograph at the top of the sheet comprises a side view of a typical...

Description of the Invention:

...0029] **Lobelia** erinus cultivar Weslospot...

...0030] Female parent: Proprietary selection of **Lobelia** erinus identified as code number 01P050, not patented
...0031] Male parent: Proprietary selection of **Lobelia** erinus identified as code number 01P508, not patented...
...0093] Plants of the new **Lobelia** have not been noted to be resistant to pathogens and pests common to **Lobelia**.

...

...0094] Plants of the new **Lobelia** have been observed to tolerate temperatures

Exemplary or Independent Claim(s):

1. A new and distinct cultivar of **Lobelia** plant named 'Weslospot', as illustrated and described.

?